

Operational Decision-Making on the Day Before and the Day of Surgery

- This talk includes many similar slides. Paging through produces animation analogous to PowerPoint animation.
- To view the slides without frustrating delays or blinks, forward from one slide to the next using either the space bar or the down arrow on your keyboard. Do not use Acrobat's Next Page navigation button, the right arrow on your keyboard, or PageDn.
- If you view the slides using View then Full Screen, first optimize your settings. Select Edit, then Preferences, then General, then Full Screen, and then No Transition.



Operational Decision-Making on the Day Before and the Day of Surgery

Franklin Dexter, M.D., Ph.D.



Director, Operations Research

Professor, Departments of Anesthesia
and Health Management & Policy

University of Iowa

Franklin-Dexter@UIowa.edu

Financial Disclosure

- Employment
 - I am employed by the University of Iowa, in part, to consult and analyze data for hospitals, anesthesia groups, and companies
 - Department of Anesthesia bills for my time
 - I receive no funds other than from the University of Iowa, including no travel reimbursement or honorarium
 - I own no healthcare stocks (other than indirectly through mutual funds)
 - I have tenure with no incentive program

Lecture 1 – Use of Ordered Priorities for Decision Making



Format of Training

- 1.5 hr to complete the scenarios
 - If you have a question not precisely matching a scenario, write it down, and wait until the end
 - Scenario 8 depends on scenario 6, which depends on scenario 4, and so forth
 - Many questions will be covered toward the end
- Record each answer and score if right/wrong
 - If “obvious,” then will get all correct



Develop Policy Guideline for Picking Up Visitor at Airport

- Spend 5 minutes to write a corporate policy for arriving at an airport sufficiently early to pick up a visiting speaker
 - Scheduled arrival time is 5 PM



Develop Policy Guideline for Picking Up Visitor at Airport

- Spend 5 minutes to write a corporate policy for arriving at an airport sufficiently early to pick up a visiting speaker
 - Scheduled arrival time is 5 PM
- Let the policy be flexible enough for workers to enjoy their work



Develop Policy Guideline for Picking Up Visitor at Airport

- Spend 5 minutes to write a corporate policy for arriving at an airport sufficiently early to pick up a visiting speaker
 - Scheduled arrival time is 5 PM
- Let the policy be flexible enough for workers to enjoy their work
 - Annette parks in Ramp A, because it is closer to the Starbuck's – she wants to get a cappuccino while waiting



Develop Policy Guideline for Picking Up Visitor at Airport

- Prevent future embarrassing events
 - Mary “borrowed” her neighbor’s sports car, and drove at an unsafe speed, reportedly to make sure that she arrived on-time
 - Frank drove at a safe speed, but arrived at 7 AM, even though the visitor was not arriving until 5 PM
 - Jim was so concerned about terrorism, that he never showed up stranding the visitor



Develop Policy Guideline for Picking Up Visitor at Airport

- Policy is that employees will make decisions in accordance with a set of ordered priorities
 - Personal preferences
 - Minimize hours showing up early
 - Safety
 - Reduce minutes that visitor has to wait
 - Show up (do not cancel)



Develop Policy Guideline for Picking Up Visitor at Airport

- Policy is that employees will make decisions in accordance with a set of ordered priorities
 - Personal preferences
 - Minimize hours showing up early
 - Safety
 - Reduce minutes that visitor has to wait
 - Show up (do not cancel)
- Order the above five priorities based on the preceding and your own scenarios



Develop Policy Guideline for Picking Up Visitor at Airport

- Policy is that employees will make decisions in accordance with a set of ordered priorities
 1. Safety
 2. Show up (do not cancel)
 3. Minimize hours showing up early
 4. Reduce minutes that visitor has to wait
 5. Personal preferences



Moral About Policies for Operational Decision-making

- Policy for operations usually results in a set of ordered priorities
- Ordered priorities cannot be chosen by consensus, because including the priorities to avoid absurd decisions results in sufficient precision that the decision is already made
- Ordered priorities cannot be chosen well in a meeting with people “thinking” about priorities



Topics of Lecture 1

- Policy on when to arrive at airport
- Decision-making based on ordered priorities
- Definitions
 - Scenarios to apply priorities
 - First-come first-served
 - Reducing tardiness, not % cases delayed
 - Bin packing



Definitions of *Staffing*, *Staff Scheduling*, and *Staff Assignment*

- Staffing (months ahead)
 - Department of Anesthesiology plans 3 ORs in the main surgical suite for urgent cases between Monday 6 PM and Tuesday 7 AM
- Staff scheduling (weeks ahead)
 - Jill is scheduled Monday 8 AM to 6 PM
 - James is scheduled Monday starting at 6 PM
- Staff assignment (0-1 days ahead)
 - Alex is assigned to OR 3



Example of *Tardiness*

- Three cases of same procedure are scheduled
- Expected OR times are 2 hr
- Expected turnover times are 30 min
- Scheduled start times: 8 AM, 10:30 AM, 1 PM
- First case entered the OR at 8:20 AM
- At 9 AM, expected *tardiness* of OR is 40 min
 - The 40 min = 20 min late + 20 min late
 - Two 20 min because there are two cases following the first case that started late



Example of *Under-Utilized OR Time*

- Staffing is planned from 7:15 AM to 3:30 PM
- An OR's last case of the day ends at 1:30 PM
- There are 2 hours of under-utilized OR time
 - Under-utilized time is from 1:30 PM to 3:30 PM



Example of *Over-Utilized OR Time*

- OR staffing is planned from 7 AM to 4 PM
- OR's last case of the day ends at 6 PM
- There are 2 hr of *over-utilized OR time*
 - Over-utilized OR time is from 4 PM to 6 PM



Precise Meaning of "Maximize OR Efficiency"

Inefficiency of use of OR time (\$) =
(Cost per hour of under-utilized OR time)
× (hours of under-utilized OR time)
+ (Cost per hour of over-utilized OR time)
× (hours of over-utilized OR time)

Strum DP et al. J Med Syst 1997



Why OR Efficiency and Not Labor Costs?



Why OR Efficiency and Not Labor Costs?

- Costs differ depending on basis for decision
 - Hospital, hospital and anesthesia providers, hospital and physicians, or society
- Costs differ depending on staff scheduling and assignment decisions
 - Even when nursing, anesthesia, and surgeons follow the same ordered-priorities, if based on labor costs they make different decisions
- Decisions based on OR efficiency invariant to these issues



Scenario 1 – Can Working Fast Increase OR Efficiency?

- OR nurses and nurse anesthetists are full-time, hourly employees
- Staffing is planned from 8 AM to 3:30 PM
- There is estimated to be 8.5 hr of cases
- Anesthesiologist gets every IV first stick, A lines and C lines first stick, and does a fiberoptic intubation in 8 minutes
- OR finishes at 3:30 PM, instead of 4:30 PM
- Has anesthesiologist increased OR efficiency?



Scenario 1 – Can Working Fast Increase OR Efficiency?

- OR nurses and nurse anesthetists are full-time, hourly employees
- On the day of surgery, the cost of an hour of under-utilized OR time is negligible relative to the cost of an hour of over-utilized OR time



Meaning of Maximizing OR Efficiency on Day of Surgery

Inefficiency of use of OR time (\$) \cong
~~(Cost per hour of under-utilized OR time)~~
× (hours of under-utilized OR time)
+ (Cost per hour of over-utilized OR time)
× (hours of over-utilized OR time)

Dexter F, Traub RD. Anesth Analg 2002

Dexter F et al. Anesthesiology 2004



Meaning of Maximizing OR Efficiency on Day of Surgery

Inefficiency of use of OR time (\$) \cong
(Cost per hour of over-utilized OR time)
 \times (hours of over-utilized OR time)



Meaning of Maximizing OR Efficiency on Day of Surgery

Inefficiency of use of OR time (\$) \cong

~~(Cost per hour of over utilized OR time)~~

\times (hours of over-utilized OR time)

Constant



Meaning of Maximizing OR Efficiency on Day of Surgery

Inefficiency of use of OR time (\$) \cong

~~(Cost per hour of over utilized OR time)~~
 \times (hours of over-utilized OR time)

Constant

- Implication
 - Maximize OR efficiency on the day of surgery by minimizing hours of over-utilized OR time



Meaning of Maximizing OR Efficiency on Day of Surgery

Inefficiency of use of OR time (\$) \cong

~~(Cost per hour of over utilized OR time)~~
 \times (hours of over-utilized OR time)

Constant

- Implication

- Maximize OR efficiency *on the day of surgery* by minimizing hours of over-utilized OR time



Scenario 1 – Can Working Fast Increase OR Efficiency?

- Scenario
 - Staffing is planned from 8 AM to 3:30 PM
 - Fast anesthesiologist finished cases in 7.5 hr instead of in the expected 8.5 hr
 - Finished at 3:30 PM instead of at 4:30 PM
 - Fast anesthesiologist increased OR efficiency by preventing 1 hr of over-utilized OR time



Scenario 1 – Can Working Fast Increase OR Efficiency?

- OR nurses, nurse anesthetists, and anesthesiologists are full-time employees
- Staffing is planned from 8 AM to ~~3:30~~ 6 PM
- There is estimated to be 8.5 hr of cases
- Anesthesiologist gets every IV first stick, A lines and C lines first stick, and does a fiberoptic intubation in 8 minutes
- OR finishes at 3:30 PM, instead of 4:30 PM
- Has anesthesiologist increased OR efficiency?



Scenario 1 – Can Working Fast Increase OR Efficiency?

- Scenario
 - Staffing is planned from 8 AM to ~~3:30~~ 6 PM
 - Fast anesthesiologist finished cases in 7.5 hr instead of in the expected 8.5 hr
 - Fast anesthesiologist ~~increased~~ *did not increase* OR efficiency



Scenario 1 – Can Working Fast Increase OR Efficiency?

- Scenario
 - Staffing is planned from 8 AM to ~~3:30~~ 6 PM
 - Fast anesthesiologist finished cases in 7.5 hr instead of in the expected 8.5 hr
 - Fast anesthesiologist ~~increased~~ *did not increase* OR efficiency

Good (rational) OR management operational decision-making is highly sensitive to the staffing for each OR, and requires knowing the staffing for each OR

Organizational Decision Making by Ordered Priorities

- Listed in order of priority
 1. Patient safety is preeminent
 2. Every surgeon has open access to OR time on *Any* future *Workday* for elective cases
 3. Maximize OR efficiency by minimizing hours of over-utilized OR time
 4. Reducing patient waiting by reducing expected tardiness for elective cases and waiting for urgent cases
 5. Personal satisfaction



Scenario 2 – Anesthesiologist Reduces Turnover Times

- Staffing is planned from 7:15 AM to 3:30 PM
- Anesthesiologist is assigned to supervise resident physicians in OR 3 and OR 4
- These ORs have just finished their first cases
- The second and last case of the day in OR 3 is expected to be finished at 2:30 PM
- The second and last case of the day in OR 4 is expected to be finished at 4:30 PM
- Which OR should anesthesiologist start next?



Scenario 2 – Anesthesiologist Reduces Turnover Times

- *Patient safety* is unaffected by decision
- Open *access* is unaffected by the decision
- *OR efficiency*
 - OR 3 expected 0 hr of over-utilized OR time
 - Finish 2:30 PM, but staffing to 3:30 PM
 - OR 4 expected 1 hr of over-utilized OR time
 - Finish 4:30 PM, but staffing to 3:30 PM
- If the patient for OR 4 is ready, the anesthesiologist should start OR 4 first



Scenario 2 – Anesthesiologist Reduces Turnover Times

- Staffing is planned from 7:15 AM to ~~3:30~~ **6 PM**
- Anesthesiologist is assigned to supervise resident physicians in OR 3 and OR 4
- These ORs have just finished their first cases
- The second and last case of the day in OR 3 is expected to be finished at 2:30 PM
- The second and last case of the day in OR 4 is expected to be finished at 4:30 PM
- Which OR should anesthesiologist start next?



Scenario 2 – Anesthesiologist Reduces Turnover Times

- *Patient safety* is unaffected decision
- Open *access* is unaffected by decision
- *OR efficiency* is unaffected by decision
 - OR 1 expected 0 over-utilized hours
 - OR 2 expected ~~1~~ 0 over-utilized hours
- *Patient waiting* is unaffected by decision
 - Last case of the day in both ORs
- *Personal satisfaction* may be affected
 - Whatever anesthesiologist thinks best



Scenario 2 – Anesthesiologist Reduces Turnover Times

- Moral
 - To make good (rational) OR management operational decisions, you need to know the staffing planned for each OR



Scenario 3 – Which OR Should Housekeeper Clean First?

- Right when two ORs are finishing their first cases of the day on time, and only one person is free to clean the two ORs
- Last case of the day in OR 12 is expected to end at 2 PM
- Last case of the day in OR 14 is expected to end at 4:30 PM
- Staffed hours are from 7 AM to 3:30 PM
- Which OR should housekeeper clean first?
 - Follow the ordered priorities



Scenario 3 – Which OR Should Housekeeper Clean First?

- *Patient safety* is unaffected by the decision
- Open *access* is unaffected by the decision
- *OR efficiency* is affected by the decision
 - OR 12 expected 0 hr of over-utilized OR time
 - Finish 2 PM, while plan staffing to 3:30 PM
 - OR 14 expected 1 hr of over-utilized OR time
 - Finish 4:30 PM, while plan staffing to 3:30 PM
- Cleaning OR 14 first is likely to increase OR efficiency



Scenario 3 – Which OR Should Housekeeper Clean First?

- Right when two ORs are finishing their first cases of the day on time, and only one person is free to clean the two ORs
- Last case of the day in OR 12 is expected to end at 2 PM
- Last case of the day in OR 14 is expected to end at 4:30 PM
- Staffing is planned from 7 AM to ~~3.30~~ 6 PM
- Which OR should housekeeper clean first?



Scenario 3 – Which OR Should Housekeeper Clean First?

- *Patient safety* is unaffected by the decision
- Open *access* is unaffected by the decision
- *OR efficiency* is unaffected by the decision
 - OR 12 expected 0 over-utilized hours
 - OR 14 expected ~~1.5~~ *0* over-utilized hours
- *Patient waiting* is unaffected by the decision
 - Last case of the day in each OR
- *Personal satisfaction* is basis for decision



Scenario 3 – Which OR Should Housekeeper Clean First?

- Why minimizing hours of over-utilized OR time, instead of minimizing overtime?



Scenario 3 – Which OR Should Housekeeper Clean First?

- Why minimizing hours of over-utilized OR time, instead of minimizing overtime?
- Same decision made regardless of ...
 - Housekeeper's scheduled hours that day
 - Housekeeper's total hours for the week
 - Collective bargaining agreements
 - How nursing schedules and assigns staff
 - How anesthesia schedules and assigns staff



Scenario 4 – Calling For the Next Patient

- Two ORs call for their next cases, but only one person is free to prepare the patients
- Both ORs are 10 min behind schedule
- The four remaining cases in OR-A are estimated to end at 2 PM
- The one remaining case in OR-B is estimated to end at 4 PM
- Staffing is planned from 7 AM to 6 PM
- Prepare which patient first?



Scenario 4 – Calling For the Next Patient

- *Patient safety* is unaffected by the decision
- Open *access* is unaffected by the decision
- *OR efficiency* is unaffected by the decision
 - OR-A expected 0 over-utilized hours
 - OR-B expected 0 over-utilized hours
- *Patient waiting* is affected by decision
 - OR-A expected total tardiness is 40 min
 - OR-B expected total tardiness is 10 min
- Prepare the patient for OR-A first



Scenario 4 – Calling For the Next Patient

- Two ORs call for their next cases, but only one person is free to prepare the patients
- Both ORs are currently on schedule
- The four remaining cases in OR-A are estimated to end at 2 PM
- The one remaining case in OR-B is estimated to end at 4 PM
- Staffing is planned from 7 AM to ~~6 PM~~ **3 PM**
- Prepare which patient first?



Scenario 4 – Calling For the Next Patient

- *Patient safety* is unaffected by the decision
- Open *access* is unaffected by the decision
- *OR efficiency* is affected by the decision
 - OR-A expected 0 over-utilized hours
 - OR-B expected 1 over-utilized hours
- Prepare the patient for OR-B first



Scenario 4 – Calling For the Next Patient

- *Patient safety* is unaffected by the decision
- Open *access* is unaffected by the decision
- *OR efficiency* is affected by the decision
 - OR-A expected 0 over-utilized hours
 - OR-B expected 1 over-utilized hours
- Prepare the patient for OR-B first

Good (rational) OR management operational decision-making is highly sensitive to the staffing planned for each OR, and requires knowing the staffing planned for each OR

Scenario 5 – Selecting Patient to Start Recovery in OR

- Staffing is planned from 7:15 AM to 3:30 PM
- At 2:20 PM, OR 11 reports that they will bring their last patient of day to PACU in 10 min
 - OR 11 has no additional cases to be performed
- At 2:25 PM, OR 21 wants to bring their 2nd to last patient of the day to the PACU in 10 min
- The PACU can currently only accept 1 extra patient while satisfying ASPAN standards
- What should PACU clerk or nurses do?



Scenario 5 – Selecting Patient to Start Recovery in OR

- *Patient safety* is unaffected by the decision
- Open *access* is unaffected by the decision
- *OR efficiency* likely is affected by the decision
 - Depending on the duration of the last case in OR 21, there may be over-utilized OR time
- *Patient waiting* is affected by the decision
 - Contact OR 11 and have them start recovering patient in the OR



Scenario 5 – Selecting Patient to Start Recovery in OR

- If months ahead plan staffing using good statistical forecasting, then at end of day there will virtually always be multiple ORs finishing last cases of day with some remaining under-utilized OR time
- If concerned about financial impact on anesthesia providers, address this tactical (not operational) issue when hospital and group creates its support agreement



Scenario 6 – Decision Based on Patient Safety

- Staffing is planned from 7:15 AM to 3:30 PM
- At 12:15, one OR is finished its elective cases
- All other ORs expected to be busy until 5 PM
- Transplant surgeon called three hours ago for cadaveric kidney transplant
 - Surgeon's assessment is that the remaining safe cold ischemic time is 10 hr
- Vascular surgery patient has ruptured AAA
- How sequence the cases based on the ordered priorities?



Scenario 6 – Decision Based on Patient Safety

- *Patient safety* is affected by decision
 - Ruptured AAA is done first
- Surgeon knows date and time of occurrence of event that requires urgent surgery or equivalently when symptoms began
 - Other examples are fracture for hip or right lower quadrant pain for appendectomy
- Hours from randomized/observational studies

Dexter F et al. J Clin Monit Comput 1999

Charalambous CP et al. Injury 2005



Scenario 7 – Moving Cases

- Staffing is planned from 7:15 AM to 3:30 PM
- OR 1 finishes last case of the day at 1:30 PM
- OR 2 is running behind
 - Its last case, scheduled from 2 PM to 3:30 PM, will not start until 5 PM
 - Anesthesia and nursing team assigned to OR 1 can perform the case safely
 - Surgeon and patient are ready
- Move the last case in OR 2 into OR 1?



Scenario 7 – Moving Cases

- *Patient safety* is unaffected by the decision
- Open *access* is unaffected by the decision
- *OR efficiency* is affected by the decision
 - Case performed entirely in over-utilized OR time if case is not moved
 - Over-utilized OR time likely reduced by at least 1.5 hr if case is moved
 - Move the case from OR 2 to OR 1



Scenario 7 – Moving Cases

- Exercise: consider the following questions
 - At start of lecture, did you specify how to pickup person at airport by using ordered priorities?
 - Would your organization hold a meeting about moving cases, brainstorm about key factors, and try to gain consensus?



Scenario 7 – Moving Cases

- Feedback from lecture is often that the use of ordered priorities is obvious
- If you answered Yes to one of the two questions, then No – not obvious



Scenario 8 – Teams Staying Late for Add-on Cases

- Hospital has one team to work past 3 PM
 - Team will not finish its case until 10 PM
- At 3 PM, all other ORs have recently finished
- There are two other 1.5 hr cases to be done, neither of which can medically wait until 10 PM, but both of which can wait to 6 PM
- What decision should be made?
 - Have one team stay late to do two cases sequentially, or have two teams stay late, each to do one case?



Scenario 8 – Teams Staying Late for Add-on Cases

- *Patient safety* is unaffected by the decision
 - Have to do the cases
- Open *access* is unaffected by the decision
 - Irrelevant since must care for the patients
- *OR efficiency* is unaffected by the decision
 - Perhaps slightly more over-utilized OR time by doing cases sequentially from the turnover
- Patient waiting is basis for the decision
 - Two teams stay late to do the cases



But, ... Personal Satisfaction is Important for Retention ...

- Why must Personal Satisfaction have a lower priority than reducing patient and surgeon waiting?



But, ... Personal Satisfaction is Important for Retention ...

- OR with 4 hr of under-utilized OR time
 - Anesthesiologist leaves for a 2 hr lunch break in between the surgeon's two cases
 - His personal satisfaction is higher priority than patient and surgeon waiting



But, ... Personal Satisfaction is Important for Retention ...

- OR with 4 hr of under-utilized OR time
 - Anesthesiologist leaves for a 2 hr lunch break in between the surgeon's two cases
 - His personal satisfaction is higher priority than patient and surgeon waiting
- **Priorities apply to operational decision-making**
 - On long term basis, if have sufficient staffing, rarely must surgeons and patients wait
 - Decisions are then usually made based on personal satisfaction



Scenario 9 – Reducing Staffing for Satisfaction

- Planned staffing is 3 ORs from 6 PM to 7 AM for non-elective cases
- Two ORs expected to finish cases at 12 MN
- At 9:30 PM, third OR is finishing its case
- Third OR covered by backup anesthesiologist (clinical next day) and junior resident
- Sole pending case can wait safely > 8 hr
- Should backup anesthesiologist stay for case?



Scenario 9 – Reducing Staffing for Satisfaction

- Over-utilized OR time
 - Expect 0 hr of over-utilized OR time regardless of when start next case, unless were very long
- Patient and surgeon waiting
 - Reduced by starting case promptly
 - Start the case now
- When and how should personal satisfaction of backup anesthesiologist be addressed?



Scenario 9 – Reducing Staffing for Satisfaction

- Personal satisfaction of backup anesthesiologist is affected by anesthesia department's staff scheduling and assignment
 - Address such issues with Vice Chair of Clinical Affairs or equivalent at appropriate time
- Staffing (i.e., OR allocations) is calculated from expected workload months in advance
 - Nursing, Anesthesia, and Surgical departments can then plan their staff scheduling and assignments independently



Scenario 10 – When Cases Exceed Staffing

- Saturday and Sunday staffing is 3 ORs x 24 hr for non-elective cases
- Not once in years have 3 ORs run non-stop for 24 hr on either Saturday or Sunday
- Under what circumstances would a 4th OR be opened on a Saturday?
 - Base list on the ordered priorities



Scenario 10 – When Cases Exceed Staffing

- *Patient safety*
 - With three ORs, a case could not reliably start by when the surgeon says it needs to start
- Open *access* to OR time
 - No effect, since do all the cases
- *OR efficiency*
 - Never open 4th OR other than for safety reasons, because would first fully fill the 3 ORs, which has never happened



Scenario 10 – When Cases Exceed Staffing

- *Patient safety*
 - With three ORs, a case could not reliably start by when the surgeon says it needs to start
 - Open *access* to OR time
 - No effect, since do all the cases
 - *OR efficiency*
 - Never open 4th OR other than for safety reasons, because would first fully fill the 3 ORs, which has never happened
- **Staffing ≠ maximum # ORs at same time**

Scenario 11 – Change in Patient Condition

- Staffing is 3 ORs from 6 PM to 7 AM
 - Two ORs will not finish cases until after 11 PM
- At 5:45 PM, two other ORs just finished cases
- At 6:00 PM, transporter leaves to pickup patient to undergo repair of elbow fracture
 - Medical deadline is to start case within 24 hr
- At 6:05 PM, OR notified of patient with expanding pseudoaneurysm of femoral artery
- Start both cases, or wait on elbow fracture case?



Scenario 11 – Change in Patient Condition

- *Patient safety* is unaffected by decision
- Open *access* to OR time unaffected by decision
 - Will do both cases
- *OR efficiency* is affected by decision
 - If do both cases right-away, elbow fracture case would be result in over-utilized OR time
 - Only start the pseudoaneurysm case



Scenario 11 – Change in Patient Condition

- Reevaluate decision whenever ...
 - New case is scheduled
 - OR times are updated
 - On-going or pending cases
 - Condition of patient waiting changes



Scenario 11 – Change in Patient Condition

- Staffing is 3 ORs from 6 PM to 7 AM
 - Two ORs will not finish cases until after 11 PM
- At 5:45 PM, two other ORs just finished cases
- At 6:00 PM, transporter leaves to pickup patient to undergo repair of elbow fracture
 - Medical deadline is to start case within 24 hr
- At 6:05 PM, OR notified of patient with expanding pseudoaneurysm of femoral artery
- At 6:10 PM, notify orthopedic resident of delay



Scenario 11 – Change in Patient Condition

- At 6:15 PM, pseudoaneurysm case starts
- At 6:15 PM, orthopedic attending updates medical deadline for case to start within 1 hr
 - Anesthesiologist's judgment is that explanation is inconsistent with evidence-based medical practice
- Unclear if start case for elbow fracture too
 - Unclear how and when address attending surgeon's basis for changing medical deadline



Scenario 11 – Change in Patient Condition

- *Patient safety* is basis for decision
 - My opinion
 - Start elbow fracture case promptly
 - OR team stays late as over-utilized OR time
- Next day, anesthesiologist reviews case with Department of Anesthesia's Vice Chair of Clinical Affairs (or equivalent person)
 - Basis for complaint is that decision was not made based on institution's ordered priorities



Three More Concepts to be Learned in Lecture 1

- First-come first-served rarely can be applied systematically in hospitals
- Why reduce tardiness as end-point for patient waiting on day of surgery?
- Principles of bin packing add-on cases

Dexter F et al. Anesthesiology 2004



First-Come First-Served at Surgical Suites

- Staffing is planned from 7 AM to 3:30 PM
- At 7 AM, urgent case has to start right away
 - Add-on case plus turnover will take 1.5 hr
- Case can be done safely in any OR
- Last cases of day will likely end at 1 PM for OR 1 and OR 2, and 3:30 PM in other ORs
- OR 1 has four short otolaryngology cases
- OR 2 has one long neurosurgery case
- Do the case in which OR?



First-Come First-Served at Surgical Suites

- OR 1 or OR 2 to minimize over-utilized hours
- OR 1: total tardiness of 6.0 hr
 - 4 patients each waiting 1.5 hr
- OR 2: total tardiness of 1.5 hr
 - 1 patient waiting 1.5 hr
- Perform urgent case in OR 2



First-Come First-Served at Surgical Suites

- OR 1 or OR 2 to minimize over-utilized hours
- OR 1: total tardiness of 6.0 hr
 - 4 patients each waiting 1.5 hr
- OR 2: total tardiness of 1.5 hr
 - 1 patient waiting 1.5 hr
- Perform urgent case in OR 2
- Cannot have answered question by first-come first-served unless relied on knowing when the elective cases were originally scheduled



First-Come First-Served at Surgical Suites

- Be on the look-out for people applying first-come first-served incorrectly
- In healthcare virtually never do you have one group of patients, and thus rarely can first-come first-served be applied rationally



Three More Concepts to be Learned in Lecture 1

- First-come first-served rarely can be applied systematically in hospitals
- Why reduce tardiness as end-point for patient waiting on day of surgery?
- Principles of bin packing add-on cases

Wachtel RE, Dexter F. Anesth Analg 2009



Reduce Percentage of Cases with Delayed Start Time?

- Staffing is planned from 7 AM to 3:30 PM
- At 7 AM, urgent case has to start right away
- OR 1 has four short otolaryngology cases
- OR 2 has one long neurosurgery case
- Hold OR 1 and OR 2 until 7:10 as decide
 - Regardless of whether case into OR 1 or OR 2, all cases are expected to have some tardiness
- Which ordered priority would be basis for decision if aim to reduce % cases delayed?



Reduce Percentage of Cases with Delayed Start Time?

- *Patient safety*
 - Start the case right away
- Open *access* is unaffected by the decision
- *OR efficiency*
 - Choose OR 1 or OR 2
- *Patient waiting* is unaffected by the decision
 - All (100%) of cases expected to be delayed regardless of the decision
- *Personal satisfaction* is basis for decision



Reduce Percentage of Cases with Delayed Start Time?

- Scenarios can be created with equivalent unusual solutions by having any percentage threshold for delay
- Reducing tardiness has advantage of not ignoring cases that have tardiness already exceeding threshold



Three More Concepts to be Learned in Lecture 1

- First-come first-served rarely can be applied systematically in hospitals
- Why reduce tardiness as end-point for patient waiting on day of surgery?
- Principles of bin packing add-on cases

Dexter F et al. Anesthesiology 1999

Dexter F, Traub RD. Anesth Analg 2002

Dexter F et al. Anesthesiology 2004



Bin Packing Surgical Cases

- Staffing planned from 7 AM to 5 PM
- Time remaining in ORs at 2 PM
 - 3 hours in OR 1 [available immediately]
 - 2 hours in OR 2 [available in 1 hr]
 - 1 hour in OR 3 [available in 2 hr]
 - 0 hours in all other ORs
- Three add-on cases listed in sequence of submission: 0.7 hr, 2.9 hr, 1.8 hr
- All can safely wait a few hours
- Perform cases in what sequence?



Bin Packing Surgical Cases

- Sort the cases based on scheduled duration from longest to shortest
 - Consider the cases in this descending order
 - Longest add-on case is assigned first
- Assign each case to OR meeting three criteria
 - Has no restrictions on equipment or personnel preventing the case from being put into the OR
 - Sufficient extra time available for the new case
 - Has least amount of additional time available

Bin Packing Surgical Cases

- Sort the cases based on scheduled duration from longest to shortest
 - Consider the cases in this descending order
 - Longest add-on case is assigned first
- Assign each case to OR meeting three criteria
 - Has no restrictions on equipment or personnel preventing the case from being put into the OR
 - Sufficient extra time available for the new case
 - Has least amount of additional time available

Safety



Bin Packing Surgical Cases

- Sort the cases based on scheduled duration from longest to shortest
 - Consider the cases in this descending order
 - Longest add-on case is assigned first
- Assign each case to OR meeting three criteria
 - Has no restrictions on equipment or personnel preventing the case from being put into the OR
 - Sufficient extra time available for the new case
 - Has least amount of additional time available

Why?



Reason for Add-on Surgical Case Scheduling Result

- On average, only 1/5 ORs has extra OR time available for add-on case
- Average time remaining in each OR each day is 1.3 hr, with large SD 1.6 hr
- Average OR time of add-on cases including their turnover times is 3.4 hr (SD 1.7 hr)
 - Add-on case scheduling applies to hospitals, since outpatient facilities rarely have add-ons



Reason for Add-on Surgical Case Scheduling Result

- Sort the cases based on scheduled duration from longest to shortest
 - Consider the cases in this descending order
 - Longest add-on case is assigned first
- Assign each case to OR meeting three criteria
 - Has no restrictions on equipment or personnel preventing the case from being put into the OR
 - Sufficient extra time available for the new case
 - Has least amount of additional time available

Why?



Reason for Add-on Surgical Case Scheduling Result

Because 0 or 1 add-on cases per OR

- Sufficient extra time available for the new case
- Has least amount of additional time available



Reason for Add-on Surgical Case Scheduling Result

- Example of importance of 0 or 1 cases
 - OR #1 has 4 hr open, OR #2 has 3 hr open
 - Add-on cases' durations are 3 hr, 2 hr, 2 hr
- If consider cases in descending sequence, and put into OR with the most remaining time
 - First, 3 hr case into OR #1
 - Second, 2 hr case into OR #2
 - Third, 2 hr case into either OR #1 or OR #2
 - Result is 1 hr over-utilized OR time, even though 0 hr was possible



Review of Topics of Lecture 1

- Policy on when to arrive at airport
- Decision-making based on ordered priorities
- Definitions
- Scenarios to apply priorities
- First-come first-served
- Reducing tardiness, not % cases delayed
- Bin packing



Lecture 2 – Impact of Uncertainties in Case Durations on Decision Making



Picking Someone Up from Airport with Uncertain Times

- Started Lecture 1 by writing policy manual for planning enough time to drive to airport to pick up a visitor
- Start this lecture by writing down appropriate percentage chance that you might have to wait in the airport for the visitor versus the visitor might have to wait for you
 - For example, write down 100% if there is certainty that you will arrive very early



Picking Someone Up from Airport with Uncertain Times

- Apply your percentage to the following driving time estimates
 - 20 min, 0% chance you wait
 - 35 min, 50% chance you wait
 - 50 min (35 min + 15 min extra “in case”)
 - 3 hr (accident?), 100% chance you wait
- Compare your time estimate to that of others



Picking Someone Up from Airport with Uncertain Times

- To improve consistency of decision-making among individuals, which of the following interventions is more important?
 - Provide everyone with historical data on all previous drives to the airport
 - Tell people what percentage value to use



Picking Someone Up from Airport with Uncertain Times

- To improve consistency of decision-making among individuals, which of the following interventions is more important?
 - Provide everyone with historical data on all previous drives to the airport
 - Tell people what percentage value to use



Moral About Data and Information Systems

- For operational decision-making, more data often will not change decision-making
- Need to learn whether the factor limiting the quality of the decision is lack of data or inconsistency in how the decision is made
 - When the primary problem is the latter, there will be lots of “who’s to blame” after software installation



How Would Complete Data on Driving Times be Used?

- If data on past 25 drives to the airport were collected by the management information system, what statistic likely would be provided to the decision-maker?
- What statistic would be ideal for driving time?



How Would Complete Data on Driving Times be Used?

- If data on past 25 drives to the airport were collected by the management information system, what statistic likely would be provided to the decision-maker?
- What statistic would be ideal for driving time?
 - Most hospital reports provide the means
 - Ideal statistic would relate to the percentage chance of being late in picking up visitor
 - For single visitor, closely related to tardiness



General Lesson About Information Systems and Data

- *If* decision is best made relying on “shortest possible” (e.g., 5th percentile) or “longest possible” (e.g., 95th percentile)
- *And If* management information system reports provides mean of historical times
- *Then*, decision using the data will likely be worse than that based on experience



General Lesson About Information Systems and Data

- *If* decision is best made relying on “shortest possible” (e.g., 5th percentile) or “longest possible” (e.g., 95th percentile)
- *And If* management information system reports provides mean of historical times
- *Then*, decision using the data will likely be worse than that based on experience

Data driven operational decision making is often no better than non-data driven decisions



Topics Covered in Lecture 2

- Picking someone up from airport (again)
- Case study on frustration with IT solution
- Applying methods based on mean duration
 - Scheduling add-on cases, moving cases, and assigning staff
- Sequencing cases to increase OR efficiency
- Upper prediction bounds to reduce waiting
- Lower prediction bounds to reduce waiting
- Time remaining in cases



Case Duration Prediction Accuracy Results

- To address complaints about surgeons waiting for other surgeons' late running cases, a hospital invests \$1.2 million in a new OR information system that tracks case durations
- Six months after implementing the new information system, and using its calculated estimated durations, the mean absolute error is *less* accurate by a small amount (5 min)
- What five reasons can explain this finding?



Case Duration Prediction Accuracy Results

- ❖ **Patient waiting does not relate to the mean historical case duration**
 - Should combine human & data information
 - Mean is relatively insensitive to more data
 - There is much process variability around the expected (mean) duration once the best method of prediction has been applied
 - There are few historical data available on which to apply statistical methods



Case Duration Prediction Accuracy Results

- ❖ **Patient waiting does not relate to the mean historical case duration**
- Should combine human & data information
 - Mean is relatively insensitive to more data
 - There is much process variability around the expected (mean) duration once the best method of prediction has been applied
 - There are few historical data available on which to apply statistical methods



Should Combine Human & Data Information

- Who will be most accurate at predicting how long it will take to drive to the airport?
 - Jim uses mean of previous times to drive
 - Jill guesses based on her experience
 - Frank uses a two-step process:
 - 1st relies on the mean of all the times that he previously drove to the airport
 - 2nd revises the estimate up by 20%, because he will drive during rush hour, even though he usually does not



Should Combine Human & Data Information

- When surgeon is provided with her historical duration, and then updates that time based on her knowledge about the case (e.g., at intra-operative briefing), the accuracy is consistently equal to or better than:
 - surgeon only estimate
 - computer only estimate

Wright IH et al. Anesthesiology 1996

Dexter F, Ledolter J. Anesthesiology 2005

Eijkemans MJC et al. Anesthesiology 2010

Dexter EU et al. Anesth Analg 2010



Case Duration Prediction Accuracy Results

- ❖ **Patient waiting does not relate to the mean historical case duration**
 - Should combine human & data information
 - Mean is relatively insensitive to more data
 - There is much process variability around the expected (mean) duration once the best method of prediction has been applied
 - There are few historical data available on which to apply statistical methods



Mean is Relatively Insensitive to More Data

- Case durations listed sequentially for one surgeon performing total hip replacement

1) 3.62 hr

2) 2.77 hr

3) 4.38 hr

4) 4.00 hr

5) 3.15 hr

6) 3.42 hr

7) 4.38 hr

8) 3.33 hr

9) 3.77 hr



Mean is Relatively Insensitive to More Data

- Case durations listed sequentially for one surgeon performing total hip replacement

1) 3.62 hr

2) 2.77 hr

3) 4.38 hr

4) 4.00 hr

5) 3.15 hr

6) 3.42 hr

7) 4.38 hr

8) 3.33 hr

9) 3.77 hr

I successively calculated mean as I added more historical data



Mean is Relatively Insensitive to More Data

- Case durations listed sequentially for one surgeon performing total hip replacement

1) 3.62 hr	3.62 hr
2) 2.77 hr	3.20 hr
3) 4.38 hr	3.59 hr
4) 4.00 hr	3.69 hr
5) 3.15 hr	3.58 hr
6) 3.42 hr	3.56 hr
7) 4.38 hr	3.67 hr
8) 3.33 hr	3.63 hr
9) 3.77 hr	3.65 hr

Final value for comparison



Mean is Relatively Insensitive to More Data

- Case durations listed sequentially for one surgeon performing total hip replacement

1) 3.62 hr 3.62 hr

2) 2.77 hr 3.20 hr

3) 4.38 hr 3.59 hr

4) 4.00 hr 3.69 hr

5) 3.15 hr 3.58 hr

6) 3.42 hr 3.56 hr

7) 4.38 hr 3.67 hr

8) 3.33 hr 3.63 hr

9) 3.77 hr 3.65 hr



Mean is Relatively Insensitive to More Data

- Incremental reduction in error in predicting mean case duration small once ≥ 3 previous cases of the same procedure(s) and surgeon
- Even if number of previous cases available to estimate case durations were increased from 1 to 39, the average tardiness would be reduced only by 2 min at a studied ambulatory surgery center and 4 min at a studied hospital

Zhou J et al. J Clin Anesth 1999



Case Duration Prediction Accuracy Results

- ❖ **Patient waiting does not relate to the mean historical case duration**
 - Should combine human & data information
 - Mean is relatively insensitive to more data
 - There is much process variability around the expected (mean) duration once the best method of prediction has been applied
 - There are few historical data available on which to apply statistical methods



Large Process Variability About Expected (Mean) Duration

- Case durations listed sequentially for one surgeon performing total hip replacement

1)	3.62 hr	-0.03 hr
2)	2.77 hr	-0.88 hr
3)	4.38 hr	0.73 hr
4)	4.00 hr	0.35 hr
5)	3.15 hr	-0.40 hr
6)	3.42 hr	-0.23 hr
7)	4.38 hr	0.73 hr
8)	3.33 hr	-0.32 hr
9)	3.77 hr	0.12 hr

Difference from mean of 3.65 hr

Large Process Variability About Expected (Mean) Duration

- Case durations listed sequentially for one surgeon performing total hip replacement

1)	3.62 hr	-0.03 hr
2)	2.77 hr	-0.88 hr
3)	4.38 hr	0.73 hr
4)	4.00 hr	0.35 hr
5)	3.15 hr	-0.40 hr
6)	3.42 hr	-0.23 hr
7)	4.38 hr	0.73 hr
8)	3.33 hr	-0.32 hr
9)	3.77 hr	0.12 hr



Large Process Variability About Expected (Mean) Duration

- Case durations listed sequentially for one surgeon performing *total hip replacement*

1)	3.62 hr	-0.03 hr
2)	2.77 hr	-0.88 hr
3)	4.38 hr	0.73 hr
4)	4.00 hr	0.35 hr
5)	3.15 hr	-0.40 hr
6)	3.42 hr	-0.23 hr
7)	4.38 hr	0.73 hr
8)	3.33 hr	-0.32 hr
9)	3.77 hr	0.12 hr

Results markedly underestimate the problem at a hospital surgical suite

Large Process Variability About Expected (Mean) Duration

- For total hip replacement, when scheduled procedure is total hip replacement, so will be the performed procedure



Large Process Variability About Expected (Mean) Duration

- For total hip replacement, when scheduled procedure is total hip replacement, so will be the performed procedure
- For many cases, there are frequent differences between scheduled procedures and actual procedures




Large Process Variability About Expected (Mean) Duration

- For total hip replacement, when scheduled procedure is total hip replacement, so will be the performed procedure
- For many cases, there are frequent differences between scheduled procedures and actual procedures
 - Cancer surgery
 - Major intra-abdominal procedures



Case duration data for scheduled Whipple procedures



Partial Solution to the Problem of Process Variability

- 6 hour is the mean case duration for the past ten Whipple procedures performed by Dr. Jones at Community General Hospital
- Dr. Jones expects his case tomorrow to be of average complexity
- Is 6 hours the expected (mean) duration of the next case?



Partial Solution to the Problem of Process Variability

- Take the mean of case durations for the past ten cases *scheduled* to be Whipple
 - Even include cases cancelled on the day of surgery before the patient enters the OR
- Doing this assures that the mean of historical durations is an unbiased estimator for the expected (mean) duration for future cases scheduled to be a Whipple procedure



Case Duration Prediction Accuracy Results

- ❖ **Patient waiting does not relate to the mean historical case duration**
 - Should combine human & data information
 - Mean is relatively insensitive to more data
 - There is much process variability around the expected (mean) duration once the best method of prediction has been applied
- **There are few historical data available on which to apply statistical methods**



Explain the Following Result

- SurgiServer OR information system provided estimated case durations using trimmed mean
 - Most recent cases (max 10) were taken of the same procedure performed by same surgeon
 - Shortest and longest durations were excluded
 - Mean was taken of the remaining eight cases
- Mean difference was 0.0 hr between estimates by sample mean and by trimmed mean
- Thinking about the equation, why was this true for most procedures?



Explain the Following Result

- SurgiServer OR information system provided estimated case durations using trimmed mean
 - Most recent cases (max 10) were taken of the same procedure performed by same surgeon
 - Shortest and longest durations were excluded
 - Mean was taken of the remaining eight cases
- Mean difference was 0.0 hr between estimates by sample mean and by trimmed mean
- Most cases have fewer than 10 occurrences of the same procedure(s) by the same surgeon

Few Historical Data Available on Which to Apply Statistics

- There are more than 4,000 different ICD-9-CM procedure codes



Few Historical Data Available on Which to Apply Statistics

- Characteristics of a hospital surgical suite
 - 11,579 cases → **1.6 : 1.0**
 - 5,156 scheduled procedures and combinations of procedures
 - 225 surgeons
 - 7,217 combinations of procedures & surgeon
- Characteristics of an outpatient surgical suite
 - 4,842 cases
 - 2,245 combinations of procedures & surgeon



Few Historical Data Available on Which to Apply Statistics

- Characteristics of a hospital surgical suite
 - 11,579 cases
 - 5,156 scheduled procedures and combinations of procedures
 - 225 surgeons
 - 7,217 combinations of procedures & surgeon
- Characteristics of an outpatient surgical suite
 - 4,842 cases
 - 2,245 combinations of procedures & surgeon

2.2 : 1.0



Few Historical Data Available on Which to Apply Statistics

- Hospital surgical suite's one year of data
 - Previous cases of the same combination of scheduled procedure(s) and surgeons
 - 37% of cases with 0 previous cases
 - 36% of cases with 1-5 previous cases
 - 12% of cases with > 19 previous cases

Zhou J et al. J Clin Anesth 1999



Few Historical Data Available on Which to Apply Statistics

- 36% of all cases performed per year in the US were of a procedure or combination of procedures performed on average once per facility per year

Dexter F, Macario A. Anesth Analg 2000



Summary and Lessons Learned

- ❖ **Patient waiting does not relate to the mean historical case duration**
 - Should combine human & data information
 - Mean is relatively insensitive to more data
 - There is much process variability around the expected (mean) duration once the best method of prediction has been applied
 - There are few historical data available on which to apply statistical methods



Summary of Lesson Learned

- If case durations were known precisely, for each case, only one estimate would be needed for its duration
- Since case durations are uncertain, one single number cannot be used for accurate decision-making
 - There is no: “*The* case duration”



Summary of Lesson Learned

- If case durations were known precisely, for each case, only one estimate would be needed for its duration
 - Since case durations are uncertain, one single number cannot be used for accurate decision-making
 - There is no: “*The* case duration”
- Understanding the problem ... to solutions



Topics Covered in Lecture 2

- Picking someone up from airport (again)
- Case study on frustration with IT solution
- Applying methods based on mean duration
 - Scheduling add-on cases, moving cases, and assigning staff
- Sequencing cases to increase OR efficiency
- Upper prediction bounds to reduce waiting
- Lower prediction bounds to reduce waiting
- Time remaining in cases



Estimating Expected (Mean) Case Duration

- If ≥ 1 previous case with same surgeon, scheduled procedure(s) and anesthetic,
 - Use the mean duration of those prior cases
- If not, and ≥ 1 previous case with same surgeon and scheduled procedure(s),
 - Use the mean duration of those prior cases
- If not, and ≥ 1 previous case with same scheduled procedure(s), ...
- If not, use the largest of the means of the individual component procedures, ...

Estimating Expected (Mean) Case Duration

- Predict expected (mean) time to complete a series of cases in same OR on the same day
 - Use sum of the means for each case



Estimating Expected (Mean) Case Duration

- More than 20 other methods that have been studied with a variety of statistical properties
- None has performed better than simply using the mean of historical cases classified using as much information as possible

Dexter F et al. J Clin Monit 1999

Macario A, Dexter F. Anesth Analg 1999



Estimating Expected (Mean) Case Duration

- More than 20 other methods that have been studied with a variety of statistical properties
- None has performed better than simply using the mean of historical cases classified using as much information as possible
- None better than just double checking surgeon's estimate manually, either

Dexter F, Ledolter J. Anesthesiology 2005



Topics Covered in Lecture 2

- Picking someone up from airport (again)
- Case study on frustration with IT solution
- Applying methods based on mean duration
 - Scheduling add-on cases, moving cases, and assigning staff
- Sequencing cases to increase OR efficiency
- Upper prediction bounds to reduce waiting
- Lower prediction bounds to reduce waiting
- Time remaining in cases



Scheduling Add-on Cases Using Bin Packing Methods

- Methods from Lecture 1 are applied to case durations estimated by taking means of historical data as on preceding slides
- At ambulatory surgery center with mean case durations of 1.6 hr, case duration inaccuracy resulted in 1.0 min of over-utilized OR time per day *vs.* knowing case durations perfectly
- At hospital with mean case durations of 3.6 hr, excess 5.4 min of over-utilized OR time

Explanation of Why Performance is so Good

- Staffing is planned from 7 AM to 3 PM
- OR 1 has two cases scheduled with estimated durations of 4 hr and 2.5 hr
- OR 2 has two cases scheduled with estimated durations of 2 hr and 2 hr
- Into which OR schedule another 2 hr case?

Dexter F et al. Anesthesiology 1999

Dexter F, Ledolter J. Anesthesiology 2005



Explanation of Why Performance is so Good

- Staffing is planned from 7 AM to 3 PM
- OR 1 has two cases scheduled with **6.5 hr** estimated durations of 4 hr and 2.5 hr
- OR 2 has two cases scheduled with **4.0 hr** estimated durations of 2 hr and 2 hr
- Into which OR schedule another 2 hr case?
- In this (and most) real-world situations, inaccuracy in predicting case duration is unlikely to have changed the original management *decision*



Implication of the Good Performance

- Negligible ROI from improved add-on case scheduling from methods to improve accuracy of case duration prediction
 - Closed circuit cameras
 - Real-time patient tracking systems
 - Graphical airport-style displays
 - More sophisticated statistical algorithms



Implication of the Good Performance

- Negligible ROI from improved add-on case scheduling from methods to improve accuracy of case duration prediction
 - Closed circuit cameras
 - Real-time patient tracking systems
 - Graphical airport-style displays
 - More sophisticated statistical algorithms
- Does this mean poor ROI from decision-support software to assist managers in scheduling add-on cases?



Implication of the Good Performance

- As considered in lecture 1, decision-support is needed to assist in how the expected (mean) durations are *used* in decision-making
- Benefit of information systems is not from improved case duration prediction, but in how the estimated case durations are *used*



Moving Cases

- Result is same as for add-on case scheduling
 - Benefit of information systems is not from improved case duration prediction, but in how the estimated case durations are used

Dexter F. Anesth Analg 2000

Dexter F et al. Anesthesiology 2004



Assigning Staff

- Suppose that OR 1 and OR 2 are still running at 5 PM, but all other ORs are finished
- One new OR team starts working at 5 PM
- Consider decision: which OR is relieved



Assigning Staff

- Estimate expected (mean) over-utilized OR time in OR1 and in OR 2
 - If mean historical duration of a case is 3 hr, and patient has been in an OR for 1 hr, then expected time remaining is slightly longer than 2 hr
- After make a preliminary decision, before relieving staff in planned OR, do a check to ensure that the OR to be relieved is not close to finishing
 - Walk to the OR and look in the window
 - Check anesthesia information system
 - Glance at a closed circuit camera
 - Phone call



Assigning Staff

- Excess over-utilized OR time from inaccuracy in predicting the time remaining is less than 1.4 min per OR per workday
 - Same reasons as for add-on case scheduling

Dexter F et al. Anesth Analg 1999



Topics Covered in Lecture 2

- Picking someone up from airport (again)
- Case study on frustration with IT solution
- Applying methods based on mean duration
 - Scheduling add-on cases, moving cases, and assigning staff
- Sequencing cases to increase OR efficiency
- Upper prediction bounds to reduce waiting
- Lower prediction bounds to reduce waiting
- Time remaining in cases



Sequencing Cases to Increase OR Efficiency

- Dr. A has two cases today in OR 1
 - 2 hr case with a surgical microscope
 - 6 hr case without specialized equipment
- Dr. B one case today in OR 2
 - 3 hr case without specialized equipment
- Dr. B wants to add a 2nd case for today
 - 4 hr case with the surgical microscope
- Yet, other surgeons have also requested OR time for briefer cases



Sequencing Cases to Increase OR Efficiency

- Dr. A has two cases today in OR 1
 - 1st: 2 hr case with the microscope
 - 2nd: 6 hr case without microscope
- Dr. B would have two cases in OR 2
 - 1st: 3 hr case without the microscope
 - 2nd: 4 hr case with the microscope



Sequencing Cases to Increase OR Efficiency

- Dr. A has two cases today in OR 1
 - 1st: 2 hr case with the microscope
 - 2nd: 6 hr case without microscope
- Dr. B would have two cases in OR 2
 - 1st: 3 hr case without the microscope
 - 2nd: 4 hr case with the microscope
- OK if Dr. A finishes 1st case before Dr. B
- What information do you want to know to judge whether uncertainty in the estimate of case duration prediction may affect decision?
- For which cases?



Sequencing Cases to Increase OR Efficiency

- Dr. A's 2 hr case with the surgical microscope
 - Mean, standard deviation, sample size
- Dr. B's 3 hr case without the microscope
 - Mean, standard deviation, sample size
- Standard deviations are used so that uncertainties in estimated mean durations are included in the calculations



Sequencing Cases to Increase OR Efficiency

- Dr. A's 2 hr case with the surgical microscope
 - Mean, standard deviation, sample size
- Dr. B's 3 hr case without the microscope
 - Mean, standard deviation, sample size
- Standard deviations are used so that uncertainties in estimated mean durations are included in the calculations
- From introductory statistics, what statistical test can be applied using this information?



Sequencing Cases to Increase OR Efficiency

- Determining probability that one case will last longer than another is similar to using *Student's t-test* to determine whether one mean exceeds another
 - Uncertainty is larger because two cases are being compared instead of two means
 - Probability estimated to accuracy $< 1.5\%$

Dexter F, Traub RD. Anesth Analg 2000

Dexter F, Ledolter J. Anesthesiology 2005



Sequencing Cases to Increase OR Efficiency

- Consider two scenarios, both with data based on surgeon and scheduled procedure(s)
 - Dr. A's 2 hr case with the microscope
 - 1st scenario: Mean 2 hr, stdev 0.5 hr, N = 30
 - 2nd scenario: Mean 2 hr, stdev 1.0 hr, N = 2
 - Dr. B's 3 hr case without the microscope
 - 1st scenario: Mean 3 hr, stdev 0.8 hr, N = 40
 - 2nd scenario: Mean 3 hr, stdev 1.5 hr, N = 3
- Schedule Dr. B to follow himself with his add-on case for 1st, but not 2nd, scenario

Sequencing Cases to Increase OR Efficiency

- For case sequencing decisions to reduce the impact of a constraint such as equipment, decision-support generally relies on analysis of relevant historical data
 - Real-time data acquisition is not needed for this problem, just immediate access to analyzed historical data



Topics Covered in Lecture 2

- Picking someone up from airport (again)
- Case study on frustration with IT solution
- Applying methods based on mean duration
 - Scheduling add-on cases, moving cases, and assigning staff
- Sequencing cases to increase OR efficiency
- Upper prediction bounds to reduce waiting
- Lower prediction bounds to reduce waiting
- Time remaining in cases



Prediction Bounds to Fill Holes in OR Schedule

- Definition
 - An upper prediction bound for the duration of a case is the value that will be exceeded by the next randomly selected case of the same type at the specified rate
 - There is a 10% chance that the duration of a case will be longer than its 90% prediction bound



Prediction Bounds to Fill Holes in OR Schedule

- Scenario showing why filling holes in the OR schedule reduces surgeon and patient waiting
 - Case 1 in OR 1 finishes unexpectedly at 10 AM
 - Case 2 in OR 1 cannot start until its scheduled time of 1 PM, because the surgeon is busy
 - There is a gap of 3 hr of available OR time
 - Dr. Holmes has an add-on case for today
 - Dr. Holmes is available to start until 12 noon
 - Dr. Holmes is then available tonight



Prediction Bounds to Fill Holes in OR Schedule

- The 90% prediction bound for the duration of Dr. Holmes' case is 2.3 hr
 - Case can be done in the 3 hr of open time with a low ($< 10\%$) risk of causing an increase in over-utilized OR time from delaying the start of the case to follow
- Use of upper prediction bounds permits reduction in waiting times, while assuring a low risk of over-utilized OR time



Prediction Bounds to *Create Holes* in OR Schedule

- Four OR ambulatory surgery center has no add-on cases
- In each of three of the four ORs, a surgeon has a list of cases for the day
- In OR 4, Dr. A has three cases scheduled from 7 AM to 10:30 AM followed by Dr. B with one case from 10:45 AM to 1 PM
- At 8 AM, Dr. A's first case has just started
- Updated end time for Dr. A's cases is 11:30



Prediction Bounds to Create Holes in OR Schedule

- Dr. B's patient can come later
- Dr. B would prefer to come later
- 90% prediction bound on the duration of Dr. B's case is 3 hr
- Dr. B can be given an updated start time of 12 noon without risking over-utilized OR time and thereby reduced OR efficiency

12 noon = 3 PM – 3 hr



Prediction Bounds Statistical Methods

- Data required to calculate a prediction bound are mean, standard deviation, and sample size of historical case duration data
- Upper prediction bounds for duration of the next one case are accurate to within 1%

Zhou J, Dexter F. Anesthesiology 1998

Dexter F et al. Anesth Analg 2001

Dexter F, Ledolter J. Anesthesiology 2005



Topics Covered in Lecture 2

- Picking someone up from airport (again)
- Case study on frustration with IT solution
- Applying methods based on mean duration
 - Scheduling add-on cases, moving cases, and assigning staff
- Sequencing cases to increase OR efficiency
- Upper prediction bounds to reduce waiting
- Lower prediction bounds to reduce waiting
- Time remaining in cases



Lower Prediction Bounds for When Patients are Ready

- Definition
 - There is 5% chance that duration of a case will be briefer than its 5% prediction bound
- Accuracy
 - The 5% prediction bounds calculated based on scheduled procedure(s) are accurate to within 0.3%

Dexter F, Traub RD. Anesthesiology 2000

Dexter F, Ledolter J. Anesthesiology 2005



Lower Prediction Bounds for When Patients are Ready

- If all patients ready before the OR is ready for them, patient waiting times would be long
- If all patients were ready just when their cases were scheduled to start, there would be marked under-utilized OR time
- Choice of when patients are ready is same as specifying the relative cost of patients' waiting time compared to cost of the staffs' idle time
 - Value you chose for airport at start of lecture



5% Prediction Bounds for When Patients are Ready

- $P < 0.05$ is common balance in medicine
 - Patients would arrive early enough that they wait 95% of time because OR is not ready, while staff wait 5% of time with empty ORs
- Median annual compensation in US of patients is 5% of sums of compensations for one patient, anesthesiologist, general surgeon, FTE housekeeper, and two nurses
- Measured actual % of occurrences ORs waited for patients at a hospital without a policy: 5%

Dexter F, Traub RD. Anesthesiology 2000

Wachtel RE, Dexter F. Anesth Analg 2007



Lower Prediction Bounds – Why Not Just An Hour or Two Early?

- Lower prediction bounds cannot be estimated accurately by using the expected (mean) duration and subtracting a safety factor
- At a hospital, patient being ready 2 hr before scheduled start of his/her case end yielded overall risk of OR staff waiting for patient of 5%
 - Among patients with a preceding case longer than 3.5 hr, the risk that the staff would wait for the patient was 14.3%



Lower Prediction Bounds – Why Not Just An Hour or Two Early?

- If a case has an expected (mean) duration of 1.5 hr, its 5% to 90% bounds may be 1.0 hr (1/3rd less) to 2.0 hr (1/3rd more)
 - Appropriate safety factor would be 0.5 hr
- If a case has an expected (mean) duration of 6.0 hr, its 5% to 90% bounds may be 4.0 hr (1/3rd less) to 8.0 hr (1/3rd more)
 - Appropriate safety factor would be 2.0 hr
- Need to use the lower prediction bounds



Lower Prediction Bounds Information Systems

- Information system can provide value by ...
 - Promptly communicating to decision-makers that there is a decision to be made
 - When patient calls or to call patients scheduled to have surgery later in the day
 - Providing immediate access to results of analysis of relevant historical data to assist them in making a quality decision



Topics Covered in Lecture 2

- Picking someone up from airport (again)
 - Case study on frustration with IT solution
 - Applying methods based on mean duration
 - Scheduling add-on cases, moving cases, and assigning staff
 - Sequencing cases to increase OR efficiency
 - Upper prediction bounds to reduce waiting
 - Lower prediction bounds to reduce waiting
- Time remaining in cases



Time Remaining in Cases – Hypothetical Example

Hours	Probability
1	10%
2	45%
3	25%
4	15%
5	5%



Time Remaining in Cases – Hypothetical Example

Hours	Probability	Hr into Case	Expected (Mean) Time Remaining
1	10%	0	2.6 hr
2	45%	1	1.8 hr
3	25%	2	1.6 hr
4	15%	3	1.3 hr
5	5%	4	1.0 hr



Time Remaining in Cases – Hypothetical Example

Hours	Probability	Hr into Case	Expected (Mean) Time Remaining
1	10%	0	2.6 hr
2	45%	1	1.8 hr
3	25%	2	1.6 hr
4	15%	3	1.3 hr
5	5%	4	1.0 hr

- $2.6 \text{ hr} = -0 + (1 \times 0.1 + 2 \times 0.45 + 3 \times 0.25 + 4 \times 0.15 + 5 \times 0.05) / 1.0$



Time Remaining in Cases – Hypothetical Example

Hours	Probability	Hr into Case	Expected (Mean) Time Remaining
1	10%	0	2.6 hr
2	45%	1	1.8 hr
3	25%	2	1.6 hr
4	15%	3	1.3 hr
5	5%	4	1.0 hr

- $2.6 \text{ hr} = -0 + (1 \times 0.1 + 2 \times 0.45 + 3 \times 0.25 + 4 \times 0.15 + 5 \times 0.05) / 1.0$



Time Remaining in Cases – Hypothetical Example

Hours	Probability	Hr into Case	Expected (Mean) Time Remaining
1	10%	0	2.6 hr
2	45%	1	1.8 hr
3	25%	2	1.6 hr
4	15%	3	1.3 hr
5	5%	4	1.0 hr

- $2.6 \text{ hr} = -0 + (1 \times 0.1 + 2 \times 0.45 + 3 \times 0.25 + 4 \times 0.15 + 5 \times 0.05) / 1.0$



Time Remaining in Cases – Hypothetical Example

Hours	Probability	Hr into Case	Expected (Mean) Time Remaining
1	10%	0	2.6 hr
2	45%	1	1.8 hr
3	25%	2	1.6 hr
4	15%	3	1.3 hr
5	5%	4	1.0 hr

- $2.6 \text{ hr} = -0 + (1 \times 0.1 + 2 \times 0.45 + 3 \times 0.25 + 4 \times 0.15 + 5 \times 0.05) / 1.0$



Time Remaining in Cases – Hypothetical Example

Hours	Probability	Hr into Case	Expected (Mean) Time Remaining
1	10%	0	2.6 hr
2	45%	1	1.8 hr
3	25%	2	1.6 hr
4	15%	3	1.3 hr
5	5%	4	1.0 hr

- $2.6 \text{ hr} = -0 + (1 \times 0.1 + 2 \times 0.45 + 3 \times 0.25 + 4 \times 0.15 + 5 \times 0.05) / 1.0$



Time Remaining in Cases – Hypothetical Example

Hours	Probability	Hr into Case	Expected (Mean) Time Remaining
1	10%	0	2.6 hr
2	45%	1	1.8 hr
3	25%	2	1.6 hr
4	15%	3	1.3 hr
5	5%	4	1.0 hr

- $2.6 \text{ hr} = -0 + (1 \times 0.1 + 2 \times 0.45 + 3 \times 0.25 + 4 \times 0.15 + 5 \times 0.05) / 1.0$



Time Remaining in Cases – Hypothetical Example

Hours	Probability	Hr into Case	Expected (Mean) Time Remaining
1	10%	0	2.6 hr
2	45%	1	1.8 hr
3	25%	2	1.6 hr
4	15%	3	1.3 hr
5	5%	4	1.0 hr

- $2.6 \text{ hr} = -0 + (1 \times 0.1 + 2 \times 0.45 + 3 \times 0.25 + 4 \times 0.15 + 5 \times 0.05) / 1.0$
- $1.8 \text{ hr} = -1 + (2 \times 0.45 + 3 \times 0.25 + 4 \times 0.15 + 5 \times 0.05) / 0.9$



Time Remaining in Cases – Hypothetical Example

Hours	Probability	Hr into Case	Expected (Mean) Time Remaining
1	10%	0	2.6 hr
2	45%	1	1.8 hr
3	25%	2	1.6 hr
4	15%	3	1.3 hr
5	5%	4	1.0 hr

- $2.6 \text{ hr} = -0 + (1 \times 0.1 + 2 \times 0.45 + 3 \times 0.25 + 4 \times 0.15 + 5 \times 0.05) / 1.0$
- $1.8 \text{ hr} = -1 + (2 \times 0.45 + 3 \times 0.25 + 4 \times 0.15 + 5 \times 0.05) / 0.9$
- $1.3 \text{ hr} = -3 + (4 \times 0.15 + 5 \times 0.05) / 0.2$




Time Remaining in Cases – Hypothetical Example

Hours	Probability	Hr into Case	Expected (Mean) Time Remaining
1	10%	0	2.6 hr
2	45%	1	1.8 hr
3	25%	2	1.6 hr
4	15%	3	1.3 hr
5	5%	4	1.0 hr

- $2.6 \text{ hr} = -0 + (1 \times 0.1 + 2 \times 0.45 + 3 \times 0.25 + 4 \times 0.15 + 5 \times 0.05) / 1.0$
- $1.8 \text{ hr} = -1 + (2 \times 0.45 + 3 \times 0.25 + 4 \times 0.15 + 5 \times 0.05) / 0.9$
- $1.3 \text{ hr} = -3 + (4 \times 0.15 + 5 \times 0.05) / 0.2$

➤ After 2.6 hr, there is not 0 hr expected time remaining!

Uncommon Non-Preemptive Tasks of Uncertain Durations




Uncommon Non-Preemptive Tasks of Uncertain Durations


- Why true?
 - Statistical property of two parameter log normal distributions with small variance terms
 - Patients with OR times of 2 hr were scheduled for the same procedure(s) as those taking 1 hr, but are undergoing different procedure(s)
 - Many procedures continue until surgery done and then closing and wakeup takes 20 min




Uncommon Non-Preemptive Tasks of Uncertain Durations



Uncommon Non-Preemptive Tasks of Uncertain Durations



Uncommon Non-Preemptive Tasks of Uncertain Durations



Uncommon Non-Preemptive Tasks of Uncertain Durations

ROOM 7

!Double Header!

NOW PLAYING: ESOPHAGECTOMY

NEXT UP: VATS LUNG BIOPSY

STARTING @ 3:30-4:20

Review of Topics Covered in Lecture 2

- Picking someone up from airport (again)
- Case study on frustration with IT solution
- Applying methods based on mean duration
 - Scheduling add-on cases, moving cases, and assigning staff
- Sequencing cases to increase OR efficiency
- Upper prediction bounds to reduce waiting
- Lower prediction bounds to reduce waiting
- Time remaining in cases



Additional Information on Operating Room Management

- www.FranklinDexter.net
 - Comprehensive bibliography of peer reviewed articles in operating room and anesthesia group management
 - Lectures on drug and supply costs, PACU staffing, OR allocation and staffing, anesthesia staffing, financial analysis, comparing surgical services among hospitals, and strategic decision making
 - Contact information

